

F 3141

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁶ : A23G 9/04, 9/02, 9/20</p>	<p>A2</p>	<p>(11) International Publication Number: WO 96/16557</p> <p>(43) International Publication Date: 6 June 1996 (06.06.96)</p>
<p>(21) International Application Number: PCT/GB95/02804</p> <p>(22) International Filing Date: 30 November 1995 (30.11.95)</p> <p>(30) Priority Data: 94308835.1 30 November 1994 (30.11.94) EP (34) Countries for which the regional or international application was filed: GB et al.</p> <p>(71) Applicant (for AU BB CA GB IE KE LK MN MW NZ SD SG SZ TT UG only): UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB).</p> <p>(71) Applicant (for all designated States except AU BB CA GB IE KE LK MN MW NZ SD SG SZ TT UG): UNILEVER N.V. [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).</p> <p>(72) Inventors: COX, David, Robert, Graham; 24 Alexandra Cottages, Tavistock Place, Bedford MK40 2RZ (GB). MOORE, Stephen, Raymond; 2 Wainwright Avenue, Thrapston, Northamptonshire NN14 4UH (GB).</p> <p>(74) Common Representative: UNILEVER PLC; Patent Division, Colworth House, Sharnbrook, Bedford MK44 1LQ (GB).</p>		<p>(81) Designated States: AL, AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>

(54) Title: PREPARATION OF ICE CRYSTALS SUSPENSION IN SUGAR SOLUTION AND ITS USE FOR MAKING CONFECTIONS

(57) Abstract

A suspension containing a large number of very small ice crystals in a sugar solution is obtained by mixing a concentrated sugar solution with a less concentrated sugar solution or water, both solutions having been cooled to a temperature of from just above the metastable limit temperature of the respective solution to just above the melting point of the respective solution prior to mixing.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

PREPARATION OF ICE CRYSTALS SUSPENSION
IN SUGAR SOLUTION AND ITS USE FOR MAKING CONFECTIONS

Field of the invention: The preparation of a suspension of ice crystals is a usual stage in the manufacture of ice confections such as water ices, granitas, ice cream and slushes. These confections are formed from an aqueous solution of sugars, which may include optional ingredients for example flavours, fats, milk proteins, emulsifiers, gas, colours and fruit pulp.

Background to the Invention:

It is desirable to be able to provide a suspension of ice crystals having an average crystal size which is as small as possible.

Conventionally the suspension of ice crystals is manufactured in a scraped surface heat exchanger (SSHE). This process mixes and cools the aqueous solution of sugars simultaneously. A disadvantage of this process is that there is a temperature gradient between the wall of the SSHE and the centre, thus crystallisation is localised at the wall of the SSHE. This results in uneven crystal distribution.

A further disadvantage of using a SSHE to provide the ice crystal suspension is that a SSHE is an expensive piece of equipment.

Alternatively the suspension of ice crystals may be manufactured via quiescent freezing process within, for example, a lolly mould. Crystallisation occurs at the walls of the mould with growth of large, dendrite crystals into the centre of the product.

We have now been able to devise a simple process for producing the suspension of ice crystals, whereby ice

crystals having the desirable small crystal size are produced. Furthermore, the process permits the formation of a large number of evenly distributed crystal nuclei. The process involves initially, individually cooling a first solution of concentrated sugar and a second solution of water or a less concentrated sugar solution prior to mixing the cooled solutions.

General description of the invention:

10

Accordingly the invention provides a method of preparing a suspension comprising ice crystals in a sugar solution, wherein a first solution of concentrated sugar and a second solution of less concentrated sugar or water are individually cooled to a temperature of from just above the metastable limit temperature of the respective solution to just above the melting point of the respective solution prior to mixing; providing that if one or both of the solutions is cooled to a temperature of from just above the metastable limit temperature of the solution to below the melting point of the solution, the respective solution is supercooled.

20

The present invention provides a convenient method of preparing a suspension containing a number of relatively small ice crystals. The size of the ice crystals can be reduced further if at least one of the solutions is supercooled prior to mixing.

25

The term "just above the melting point" means the temperature is in the range up to about 5°C above the melting point for ice in the solution.

30

The term "just above the metastable limit temperature" means the temperature is about 0.5°C above the metastable limit temperature for the solution or higher.

35

The term "metastable limit temperature" is a well known term within the art. A suitable method for measurement is described in "Crystallization", J Mullin, Butterworth 1972, p178 & 179 & Fig 6.4.

The first solution is a concentrated sugar solution. This solution will preferably have a concentration above about 45% w/w more preferably above about 55%.

The second solution is either water or a less concentrated sugar solution. Preferably the second solution is water because this the most efficient method for commercial use. The less concentrated sugar solution will preferably have a concentration below about 20% w/w, more preferably below about 10%.

The sugar will usually be sucrose but other sugars, eg glucose, dextrose and fructose can be used. In some procedures there ~~may be~~ a benefit in having different sugars in the two solutions.

Either solution may contain small quantities of flavouring, citric acid, sodium citrate, fats, milk proteins, emulsifiers, gas, fruit pulp, fruit juice or stabiliser.

The use of greater than two solutions having either different or identical sugar concentrations, each solution being individually cooled to a temperature of from just above the metastable limit temperature of the solution to just above the melting point of the solution prior to mixing, is clearly possible and falls within the scope of the present invention as claimed; providing that at least one solution is of concentrated sugar and at least one solution is of less concentrated sugar or water.

Typically the process of the invention will provide an ice suspension containing from 0.5 to 10% by weight ice

crystals in a sugar solution. This ice suspension forms rapidly. The level of ice formed is too low for commercial use as, for example, an ice cream or a water ice. Therefore a second freezing step is required to increase the phase volume of ice in the suspension. This second freezing step is required to increase the phase volume of ice in the suspension. This second freezing step may occur simultaneously with the mixing step. Alternatively, the ice suspension according to the invention may be prepared first and then introduced into a suitable freezing system.

Advantageously any convenient means for mixing the cooled solutions may be used. Use of different mixing processes will modify the size and/or shape of the crystals produced.

Thus static mixers which pass streams of liquid through a non-moving mechanical device arranged to bring the streams together and rapidly mix them can be used. An example is the Kenics mixer. ~~This route~~ provides a resultant stream containing the suspension of fine ice crystals. The method of the invention also extends to introducing the streams into a vessel, eg a mould, optionally containing a stirrer. This route allows the suspension to be formed in a vessel in a coolant, so the suspension is then quiescently frozen.

An alternative route is to have the mixing step performed in a vessel capable of providing dynamic freezing, eg scraped surface heat exchanger (SSHE) or a cavity transfer mixer. In this route the suspension of ice crystals is formed within the apparatus providing the second freezing step.

Thus the method forms the large number of small crystals in the bulk of the water and not on the surface of a containing vessel.

After preparing the suspension as a stream it can be led to

a SSHE within which the freezing is completed. Aeration may be introduced at this stage.

5 One or both of the solutions may be supercooled prior to mixing. Supercooling enables particularly small ice crystals to be formed.

10 By supercooled is meant that the solution has been cooled to a lower temperature than the melting temperature for ice in solution without ice being formed. In order to supercool a solution, the solution is cooled slowly. Supercooling may be achieved by any known technique or may be effected by cooling the solution in a system in which the coolant temperature is never below the metastable limit
15 of the solution.

20 In order to provide the desired final ice confection, eg water ice, ice cream, granita, slush, the resultant suspension of ice crystals may be mixed with the components of the ice confection. Alternatively, and preferably, the components of the ice confection may be added to one or both of the solutions used to prepare the initial ice suspension. Typically the additional components of the ice
25 confection will be included in the water or less concentrated sugar solution. A further alternative is the components of the ice confection may be added at the mixing stage.

30 **Specific description of the invention:**

Examples will now be given to illustrate but not limit the method of the invention.

35 Water ice slurries were formed by mixing two solutions.

A concentrated sucrose stream A (30-62% w/w sucrose) was cooled to a temperature in the range 0-4°C higher than its

equilibrium melting point. This was mixed with a dilute sucrose stream B (0-15%w/w sucrose) which was cooled to a temperature in the range 0-4°C higher than its equilibrium melting point. Either stream may contain small quantities of flavouring, citric acid, sodium citrate, fruit juice or stabilizer, eg guar and LBG. The two streams were mixed at a weight ratio between 8:1 and 1:2 of the concentrated (A) to the dilute stream (B). The total flow was 300 Kghr⁻¹. This created a mixed stream with an initial temperature lower than the equilibrium melting point of the stream and some of the water crystallised. The stream had a temperature in the range from -18°C to -3.0°C.

The mixer used was a Kenics static tube mixer (obtainable from Chemineer Ltd, Derby, England) comprising a tube 30 cm long with a 2.5 cm diameter. This tube contained 12 curved elements arranged along the tube parallel to the liquid flow with alternative elements being perpendicular to the adjacent elements. Each curved element rotates the liquid flow in a different direction.

An alternative mixer is the Sulzer type (obtainable from Sulzer-Chemtech, Winterhur, Switzerland).

This mixing created a physical environment in which ice crystals formed spontaneously. A water ice slurry was formed which contained between 0.5% and 15% w/w ice. Four examples were performed and their conditions are quoted in Tables I and II. The resultant ice slurries had calculated mass fractions from 1% to 5%.

The final structure obtained after the second freezing stage had a soft texture and was distinct from the dendritic structure provided by quiescent freezing.

Table I

Ex	STREAM A		STREAM B		Flow Ratio (A/B)	Ice content* (mass fraction)
	Sucrose concentration (%w/w)	Temperature (°C)	Sucrose concentration (%w/w)	Temperature (°C)		
I	62	-13.5	0	0	1	0.05
II	50	-7.1	0	0	1.6	0.01
III	62	-13.5	15	-1.1	0.5	0.02
IV	55	-9.2	15	-1.1	0.67	0.01

* calculated

Table II

Melting points of solutions used:

Sucrose (%w/w)°C

none

0°

15

-1°C

50

-7.5°C

55

-9.3°C

62

-13.8°C

This slurry, which comprised a mass of very small ice crystals, was then frozen further in a second stage in a blast freezer or a similar apparatus to give a water ice with an unusual structure which retained the large number of small crystals. The resultant ice slurry was subjected to downstream freezing to generate a commercially usable water ice. This downstream, ie post mixing, freezing stage can be quiescent, for example in moulds passed through freezing zones, or with movement, for example in a scraped surface heat exchanger. Examples of freezing zones for quiescent freezing are brine baths in which moulds are placed and freezing tunnels using very cold gasses. The precise freezing regime used will depend on the structure

and shape desired in the final product.

5 A process identical to the above was used to prepare aerated water ice. Either the concentrated or the dilute stream was aerated with air (or any suitable gas) prior to the mixing step. Small air bubbles were distributed throughout the suspension and became entrapped in the frozen product.

10 The method is particularly applicable to the formation of water ices but can also be used in the preparation of other foodstuffs in which ice is a component. Thus the method can be used in the preparation of ice cream.

15 Example V

A water ice slurry was formed by mixing two solutions.

20 A concentrated sucrose stream A (62 wt%) was cooled to -10.9°C . This was mixed with water having additionally 1.4 wt% citric acid, 0.7 wt% sodium citrate, 0.3 wt% lemon flavour which had been cooled to 0.3°C . The two streams were mixed at a weight ratio of 1:1. The total flow was 3.3 kg/min. The mixer used was a 12 element kenics mixer.
25 Ice crystals were observed in the exit stream which had a temperature of -2.1°C .

Example VI

30 A water ice solution was formed by mixing two solutions.

0.45 kg of 55 wt% sucrose solution (melting point -9.2°C) was cooled to -7.5°C . This was mixed with 0.5 kg of water which had been supercooled to -1.7°C . Mixing was achieved
35 using a Rushton turbine rotating at 200 rpm. Ice crystals were observed to form almost immediately in the mixture.

Example VII

5 Sucrose solution (60 wt%) at -12.8°C was mixed with water at 0.1°C in a weight ratio of 3:2 in a Rushton turbine mixer at 240 rpm to form stream A containing a slurry of ice crystals.

Stream B was a composition containing, by weight,

10	water	53.8%
	skimmed milk powder	25.0%
	butter fat	20.0%
	locust bean gum	0.4%
	emulsifier	0.8%

15

which was pasteurised and homogenised. Streams A and B (at -2.6°C) were mixed at a weight ratio of 1:1 and the resulting stream was simultaneously aerated with added Nitrogen and sheared to give a stiff product. This product
20 was then hardened quiescently to -16°C to give an ice cream confection.

Claims

1. A method of preparing a suspension comprising ice crystals in a sugar solution, wherein a first solution of concentrated sugar and a second solution of less concentrated sugar or water are individually cooled to a temperature of from just above the metastable limit temperature of the respective solution to just above the melting point of the respective solution prior to mixing; providing that if one or both of the solutions is cooled to a temperature of from just above the metastable limit temperature of the respective solution to below the melting point of the respective solution, the respective solution is supercooled.
2. A method according to claim 1 wherein the first solution is a concentrated sugar solution having a sugar concentration above about 45% w/w, preferably above about 55% w/w.
3. A method according to claim 1 or 2 wherein the second solution is water or a less concentrated sugar solution having a sugar concentration below about 20% w/w preferably below about 10% w/w.
4. A method according to any preceding claim wherein the second solution is water.
5. A method according to any preceding claim wherein at least one solution has a temperature of from just above the metastable limit temperature of the solution to below the melting point of the solution.
6. A method according to any preceding claim wherein both the first and the second solutions have a temperature of from just above the metastable limit temperature of the respective solution to below the melting point of the


respective solution.

5 7. A method according to any one of claims 1 to 4 wherein at least one solution has a temperature at or just above its melting point.

10 8. A method according to any one of claims 1 to 4 wherein both the first and the second solutions have a temperature at or just above their respective melting points.

15 9. A method of preparing a product containing an ice suspension wherein the suspension of any preceding claim is subjected to a freezing step simultaneous to the mixing step.

20 10. A method of preparing a product containing an ice suspension wherein the suspension of any one of claims 1 to 8 is subjected to a freezing step after mixing.



PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 :

A23G 9/04, 9/02, 9/20

A3

(11) International Publication Number:

WO 96/16557

(43) International Publication Date:

6 June 1996 (06.06.96)

(21) International Application Number: PCT/GB95/02804

(22) International Filing Date: 30 November 1995 (30.11.95)

(30) Priority Data:

94308835.1

30 November 1994 (30.11.94)

EP

(34) Countries for which the regional or
international application was filed:

GB et al.

(71) Applicant (for AU BB CA GB IE KE LK MN MW NZ SD SG SZ
TT UG only): UNILEVER PLC [GB/GB]; Unilever House,
Blackfriars, London EC4P 4BQ (GB).(71) Applicant (for all designated States except AU BB CA GB IE
KE LK MN MW NZ SD SG SZ TT UG): UNILEVER N.V.
[NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).(72) Inventors: COX, David, Robert, Graham; 24 Alexandra Cot-
tages, Tavistock Place, Bedford MK40 2RZ (GB). MOORE,
Stephen, Raymond; 2 Wainwright Avenue, Thrapston,
Northamptonshire NN14 4UH (GB).(74) Common Representative: UNILEVER PLC; Patent Division,
Colworth House, Sharnbrook, Bedford MK44 1LQ (GB).(81) Designated States: AL, AM, AT, AU, BB, BG, BR, BY, CA,
CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP,
KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD,
MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,
SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European
patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE,
LS, MW, SD, SZ, UG).**Published***With international search report.*(88) Date of publication of the international search report:
29 August 1996 (29.08.96)

(54) Title: PREPARATION OF ICE CRYSTALS SUSPENSION IN SUGAR SOLUTION AND ITS USE FOR MAKING CONFECTIONS

(57) Abstract

A suspension containing a large number of very small ice crystals in a sugar solution is obtained by mixing a concentrated sugar solution with a less concentrated sugar solution or water, both solutions having been cooled to a temperature of from just above the metastable limit temperature of the respective solution to just above the melting point of the respective solution prior to mixing.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Larvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 95/02804

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A23G9/04 A23G9/02 A23G9/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A23G

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US,A,3 987 211 (L. D. DUNN) 19 October 1976 see the whole document	1,4-7,9
Y	US,A,3 343 967 (I. H. RUBENSTEIN) 26 September 1967 see the whole document	1,9,10
Y	US,A,2 941 885 (B. E. TOMINSON) 21 June 1960 see column 2, line 38 - column 3, line 48 see column 2, line 12-14; claims; figure 1	1,9,10
A	GB,A,2 251 863 (SHUICHI KARITA ETR AL.) 22 July 1992 see the whole document	1
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *&* document member of the same patent family

Date of the actual completion of the international search

4 June 1996

Date of mailing of the international search report

14.06.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Guyon, R

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 95/02804

C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US,A,3 949 102 (J.A.HELLYER ET AL.) 6 April 1976 see column 4, line 55 - column 6, line 56; claim 4; examples 4,5 ---	1
A	EP,A,0 201 141 (UNILEVER) 17 December 1986 see column 1, line 50-54; claims; figures ---	9,10
A	US,A,3 897 571 (B. HOMLER ET AL.) 29 July 1975 see column 2, line 64-66 see column 3, line 1-65 ---	1
A	FR,A,2 275 155 (UNILEVER) 16 January 1976 ---	
A	EP,A,0 168 537 (GOLDSTEIN VLAD) 22 January 1986 ---	
X	PATENT ABSTRACTS OF JAPAN vol. 13, no. 15 (C-559), 13 January 1989 & JP,A,63 222647 (YAMANE HIROSHI ET AL.), 16 September 1988, see abstract ---	1
A	US,A,4 808 428 (S. L. FORSSTROM ET AL.) 28 February 1989 ---	
A	FR,A,2 268 472 (GENERAL FOODS CORP.) 21 November 1975 -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 95/02804

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-3987211	19-10-76	NONE	
US-A-3343967	26-09-67	NONE	
US-A-2941885	21-06-60	NONE	
GB-A-2251863	22-07-92	JP-A- 4252171 HK-A- 30595 US-A- 5296251	08-09-92 17-03-95 22-03-94
US-A-3949102	06-04-76	BE-A- 843603 CA-A- 1082979 DE-A- 2629023 FR-A, B 2315857 GB-A- 1542698 JP-A- 52031868 NL-A- 7607152	30-12-76 05-08-80 03-02-77 28-01-77 21-03-79 10-03-77 03-01-77
EP-A-201141	12-11-86	AU-B- 586249 AU-B- 5723386 AU-B- 586250 AU-B- 5723486 DE-A- 3681062 JP-B- 7083682 JP-A- 61265051 PT-B- 82552 US-A- 4746523	06-07-89 18-12-86 06-07-89 13-11-86 02-10-91 13-09-95 22-11-86 31-12-93 24-05-88
US-A-3897571	29-07-75	NONE	
FR-A-2275155	16-01-76	GB-A- 1523591 AU-B- 8158375 BE-A- 829607 DE-A- 2523995 JP-A- 51007154 LU-A- 72608 NL-A- 7506492 SE-A- 7506227 US-A- 4039693	06-09-78 02-12-76 28-11-75 18-12-75 21-01-76 17-03-76 02-12-75 01-12-75 02-08-77

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 95/02804

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-168537	22-01-86	AT-T- 119656	15-03-95
		AT-T- 119658	15-03-95
		AU-B- 583051	20-04-89
		AU-B- 4722885	10-02-86
		CA-A- 1276475	20-11-90
		DE-D- 3486374	13-04-95
		DE-T- 3486374	02-11-95
		DE-D- 3486375	13-04-95
		DE-T- 3486375	02-11-95
		EP-A- 0322705	05-07-89
		EP-A- 0316966	24-05-89
		JP-T- 62500043	08-01-87
		NO-C- 163304	23-04-96
		WO-A- 8600692	30-01-86
US-A-4808428	28-02-89	AU-B- 3132889	21-09-89
		CA-A- 1334353	14-02-95
		EP-A,B 0336462	11-10-89
FR-A-2268472	21-11-75	BE-A- 828304	24-10-75
		CA-A- 1050812	20-03-79
		CH-A- 604564	15-09-78
		DE-A- 2518302	06-11-75
		GB-A- 1458934	15-12-76
		JP-C- 1167513	08-09-83
		JP-A- 51009766	26-01-76
		JP-B- 57046818	05-10-82
		NL-A- 7504851	28-10-75
		SE-A- 7504778	27-10-75